Author Index

Achtert, G., see Dahmen, W. 157
Adams, P.C.
—, Roberts, F.D., Powell, L.W. and Halliday, J.W.
Chromatographic detergent exchange in the preparation of membrane receptor protein 341
Anlér, E.-L., see Johansson, M. 55
Aronson, A.L., see Tyczkowska, K. 103
Beaudoin, N., see Dubé, L.M. 113
Becker, J.U., see Krämer, B.K. 351
Belfroid, R.D.M., see Van der Krogt, J.A. 9
Bergqvist, Y.
—, Eckerbom, S., Larsson, N. and Churchill, F.C.
High-performance liquid chromatographic separation and extraction investigation for the simultaneous determination of mefloquine and its carboxylic acid metabolite 295
Bies, J.M., see Schneebaum, S. 166
Boomsma, F.
—, Van der Hoorn, F.A.J., Man in 't Veld, A.J. and Schalekamp, M.A.D.H.
Determination of D,L-*threo*-3,4-dihydroxyphenylserine and of the D- and L-enantiomers in human plasma and urine 219
Bremm, K.D., see Knöller, J. 257
Bresson, M., see Rop, P.P. 172
Briele, H.A., see Schneebaum, S. 166
Brodfuehrer, J.I.
— and Powis, G.
Gas chromatographic assay for the new antitumor agent sulfamic acid diester (NSC 329680) and its stability in buffer, blood and plasma 247
Canfell, P.C., see Furuta, T. 41
Carlin, J.R.
—, Christofalo, P. and Vandenheuvel, W.J.A.
High-performance liquid chromatographic determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5α-androst-1-ene-17β-carboxamide, a 4-azasteroid, in human plasma from a phase I study 79
Castagnoli, K.P., see Furuta, T. 41
Chai, M., see Emm, T.A. 162
Chain, B.M., see Wallace, G.R. 239
Christofalo, P., see Carlin, J.R. 79
Churchill, F.C., see Bergqvist, Y. 295
Cooper, T.B., see Suckow, R.F. 287
Curtis, M.A., see Koenigbauer, M.J. 277
Dahmen, W.
—, Gielsdorf, W., Jeremic, B., Achtert, G. and Hausleiter, H.-J.
Determination of eproxindine hydrochloride in human plasma by capillary gas chromatography 157
De Leenheer, A.P., see Lambert, W.E. 195
DePace, A., see Verebey, K. 151
Dubé, L.M.
—, Beaudoin, N., Lalande, M. and McGilveray, I.J.
Determination of nalbuphine by high-performance liquid chromatography with electrochemical detection: application to clinical samples from post-operative patients 113
Dubois, J.P., see Godbillon, J. 269
Duval, M., see Godbillon, J. 269
Eckerbom, S., see Bergqvist, Y. 295
Emm, T.
—, Lesko, L.J., Leslie, J. and Perkal, M.B.
Determination of albuterol in human serum by reversed-phase high-performance liquid chromatography with electrochemical detection 188
—, Leslie, J., Chai, M., Lesko, L.J. and Perkal, M.B.
High-performance liquid chromatographic assay of cephalexin in serum and urine 162
Engels, W.
—, Kamps, M.A.F., Lemmens, P.J.M.R., Van der Vusse, G.J. and Reneman, R.S.
Determination of prostaglandins and thromboxane in whole blood by high-performance liquid chromatography with fluorimetric detection 209
Ennachachibi, A.
—, Nicolas, P., Fauvelle, F., Perret, G. and Petitjean, O.
Effective high-performance liquid chromatographic determination of glafenine in plasma: pharmacokinetic application 307
Evelo, C.T.A.
—, Kon, T.C., Fijneman, P.H.S. and Henderson, P.Th.
Separation of human haemoglobin alkylated at β93 cysteine from its native form by fast protein liquid chromatography 335
Fauvelle, F., see Ennachachibi, A. 307
Fijneman, P.H.S., see Evelo, C.T.A. 335

Furuta, T.
—, Canfell, P.C., Castagnoli, K.P., Sharma, M.L. and Miller, R.D.
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary gas chromatography using nitrogen-sensitive detection 41
Gauron, S., see Godbillon, J. 269
Gielsdorf, W., see Dahmen, W. 157
Godbillon, J.
—, Duval, M., Gauron, S. and Dubois, J.P.
High-performance liquid chromatographic determination of the penem antiobiotic (5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penem-3-carboxylic acid in human plasma and urine 269
Gottvaldova, M.
—, Hrabova, H., Sillinger, V. and Kucera, J.
Biospecific sorption of cyclodextrin glucosyltransferase on physically modified starch 331
Gupta, R.N.
Determination of catecholamines in body fluids 359
Hadidi, A.-H.F.A.
— and Idle, J.R.
Combined thin-layer chromatography–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 121
Halliday, J.W., see Adams, P.C. 341
Hara, S., see Ijitsu, T. 29
Hausleiter, H.-J., see Dahmen, W. 157
Heirwegh, K.P.M., see Rothuizen, J. 19
Henderson, P.Th., see Evelo, C.T.A. 335
Hensler, T., see Knöller, J. 199
Henton, D., see Vreman, H.J. 315
Hoorn, F.A.J. van der, see Boomsma, F. 219
Hoyer, G.L.
Improved high-performance liquid chromatographic method for the analysis of serum sotalol 181
Hrabova, H., see Gottvaldova, M. 331
Ibe, K., see Köppel, C. 144
Idle, J.R., see Hadidi, A.-H.F.A. 121
Ijitsu, T.
—, Ueno, M. and Hara, S.
Two-dimensional high-performance liquid chromatographic system for the direct determination of tocopherol derivatives in plasma 29
Jeremic, B., see Dahmen, W. 157
Johansson, M.
— and Anlér, E.-L.
Gas chromatographic analysis of flunixin in equine urine after extractive methylation 55
Kamata, S., see Yonekura, T. 320
Kamps, M.A.F., see Engels, W. 209
Kanfer, I., see Stubbs, C. 93
Knöller, J.
—, König, W., Schönfeld, W., Bremm, K.D. and Köller, M.
Application of high-performance liquid chromatography of some antibiotics in clinical microbiology 257
—, Schönfeld, W., Köller, M., Hensler, T. and König, W.
Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis – routine monitoring by high-performance liquid chromatography 199
Koel, M., see Nebinger, P. 326
Köller, M., see Knöller, J. 199, 257
König, W., see Knöller, J. 199, 257
Koenigbauer, M.J.
— and Curtis, M.A.
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277
Köppel, C.
—, Tenczer, J. and Ibe, K.
Gas chromatographic–mass spectrometric study of urinary metabolism of melperone 144
Kon, T.C., see Evelo, C.T.A. 335
Koppele, J.M. te, see Te Koppele, J.M. 67
Kouwen, A.M. van, see Rothuizen, J. 19
Krämer, B.K.
—, Mayer, F., Liebich, H.M., Ress, K.M., Kühlkamp, V., Becker, J.U., Risler, T. and Seipel, L.
Rapid and inexpensive high-performance liquid chromatographic method for the quantification of flecainide in serum 351
Krogt, J.A. van der, see Van der Krogt, J.A. 9
Kucera, J., see Gottvaldova, M. 331
Kühlkamp, V., see Krämer, B.K. 351
Lalande, M., see Dubé, L.M. 113
Lambert, W.E.
—, De Leenheer, A.P., Walgraeve, H.R. and Tassaneeyakul, W.
Monitoring of cyclosporin in whole blood by reversed-phase liquid chromatography on a butyl column 195

Larsson, N., see Bergqvist, Y. 295
Leenheer, A.P. De, see Lambert, W.E. 195
Lemmens, P.J.M.R., see Engels, W. 209
Lesko, L.J., see Emm, T. 162, 188
Leslie, J., see Emm, T. 162, 188
Liebich, H.M., see Krämer, B.K. 351
Lucarelli, C., see Marzo, A. 345
McGilveray, I.J., see Dubé, L.M. 113
McLeod, A., see Wallace, G.R. 239
Man in 't Veld, A.J., see Boomsma, F. 219
Mark, E.J. van der, see Te Koppele, J.M. 67
Marzo, A.
— and Lucarelli, C.
Gas chromatographic evaluation of buflomedil in biological samples employing the thermoionic specific detector 345
Mayer, F., see Krämer, B.K. 351
Mefford, I.N.
Determination of catecholamines in body fluids 360
Miller, R.D., see Furuta, T. 41
Mulder, G.J., see Te Koppele, J.M. 67
Nagatsu, T., see Naoi, M. 229
Naoi, M.
—, Takahashi, T. and Nagatsu, T.
Simple assay procedure for tyrosine hydroxylase activity by high-performance liquid chromatography employing coulometric detection with minimal sample preparation 229
Nebinger, P.
— and Koel, M.
Automatic determination of serotonin in biological fluids by liquid chromatography 326
Nicolas, P., see Ennachachibi, A. 307
Okada, A., see Yonekura, T. 320
Perkal, M.B., see Emm, T. 162, 188
Perret, G., see Ennachachibi, A. 307
Petitjean, O., see Ennachachibi, A. 307
Powell, L.W., see Adams, P.C. 341
Powis, G., see Brodfuehrer, J.I. 247
Quicke, J., see Rop, P.P. 172
Reneman, R.S., see Engels, W. 209
Ress, K.M., see Krämer, B.K. 351
Risler, T., see Krämer, B.K. 351
Roberts, F.D., see Adams, P.C. 341
Rop, P.P.
—, Spinazzola, J., Zahra, A., Bresson, M., Quicke, J. and Viala, A.
Column liquid chromatographic analysis of barbiturates in biological fluids 172
Rosenthal, P., see Vreman, H.J. 315
Rothuizen, J.
—, Heirwegh, K.P.M. and Van Kouwen, A.M.
Novel method for high-performance liquid chromatography of azo derivatives of conjugated and unconjugated bilirubin 19
Salari, H., see Steffenrud, S. 1
Sandnes, K., see Waagboe, R. 138
Schalekamp, M.A.D.H., see Boomsma, F. 219
Schneebaum, S.
—, Bies, J.M. and Briele, H.A.
Reversed-phase high-performance liquid chromatography of dactinomycin 166
Schönfeld, W., see Knöller, J. 199, 257
Seipel, L., see Krämer, B.K. 351
Sharma, M.L., see Furuta, T. 41
Sillinger, V., see Gottvaldova, M. 331
Spahn, H.
S-(+)-Flunoxaprofen chloride as chiral fluorescent reagent 131
Spinazzola, J., see Rop, P.P. 172
Steffenrud, S.
— and Salari, H.
Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 1
Stevenson, D.K., see Vreman, H.J. 315
Stubbs, C.
— and Kanfer, I.
High-performance liquid chromatography of erythromycin propionyl ester and erythromycin base in biological fluids 93
Suckow, R.F.
— and Cooper, T.B.
Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative liquid chromatography 287
Takahashi, T., see Naoi, M. 229
Tassaneeyakul, W., see Lambert, W.E. 195
Te Koppele, J.M.
—, Van der Mark, E.J. and Mulder, G.J.
Liquid chromatographic determination of diastereomeric glutathione conjugates and further derivatives of α-bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 67
Tenczer, J., see Köppel, C. 144
Tyczkowska, K.
— and Aronson, A.L.
Analysis of cephalexin from canine skin biopsy by liquid chromatography with ultraviolet–visible photodiode-array detection 103
Ueno, M., see Ijitsu, T. 29
Valkenburg, C.F.M. van, see Van der Krogt, J.A. 9
Vandenheuvel, W.J.A., see Carlin, J.R. 79
Van der Hoorn, F.A.J., see Boomsma, F. 219

Van der Krogt, J.A.
—, Van Valkenburg, C.F.M. and Belfroid, R.D.M.
Tritium isotope effect in reversed-phase high-performance liquid chromatographic analysis of dopamine and dihydroxyphenylacetic acid 9
Van der Mark, E.J., see Te Koppele, J.M. 67
Van der Vusse, G.J., see Engels, W. 209
Van Kouwen, A.M., see Rothuizen, J. 19
Van Valkenburg, C.F.M., see Van der Krogt, J.A. 9
Verebey, K.
— and DePace, A.
Rapid confirmation of enzyme-multiplied immunoassay test (EMIT) for phencyclidine-positive urine samples with capillary gas chromatography–nitrogen–phosphorus detection 151
Viala, A., see Rop, P.P. 172
Vreman, H.J.
—, Stevenson, D.K., Henton, D. and Rosenthal, P.
Correlation of carbon monoxide and bilirubin production by tissue homogenates 315
Vusse, G.J. van der, see Engels, W. 209
Waagboe, R.
— and Sandnes, K.
Determination of vitellogenin in serum of rainbow trout (*Salmo gairdneri*) by high-performance gel permeation chromatography 138
Wada, H., see Yonekura, T. 320
Walgraeve, H.R., see Lambert, W.E. 195
Wallace, G.R.
—, McLeod, A. and Chain, B.M.
Chromatographic analysis of the trinitrophenyl derivatives of insulin 239
Wasa, M., see Yonekura, T. 320
Watanabe, T., see Yonekura, T. 320
Yamatodani, A., see Yonekura, T. 320
Yonekura, T.
—, Kamata, S., Wasa, M., Okada, A., Yamatodani, A., Watanabe, T. and Wada, H.
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by high-performance liquid chromatography using derivatization with fluorescamine 320
Zahra, A., see Rop, P.P. 172

Subject Index

Acylaminopenicillins
Application of HPLC of some antibiotics in clinical microbiology 257
Albuterol
Determination of albuterol in human serum by reversed-phase HPLC with electrochemical detection 188
Amines
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 320
(5*R*,6*S*)-2-Aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penem-3-carboxylic acid
HPLC determination of the penem antibiotic (5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penem-3-carboxylic acid in human plasma and urine 269
Antibiotics
Application of HPLC of some antibiotics in clinical microbiology 257
Antibiotics
HPLC determination of the penem antibiotic (5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penem-3-carboxylic acid in human plasma and urine 269
Arachidonic acid metabolites
Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis - routine monitoring by HPLC 199
Aspoxicillin
Application of HPLC of some antibiotics in clinical microbiology 257
4-Azasteroid
HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5α-androst-1-ene-17β-carboxamide, a 4-azasteroid, in human plasma from a phase I study 79
Azlocillin
Application of HPLC of some antibiotics in clinical microbiology 257
Barbiturates
Column LC chromatographic analysis of barbiturates in biological fluids 172
Bilirubin
Novel method for HPLC of azo derivatives of conjugated and unconjugated bilirubin 19
Bilirubin
Correlation of carbon monoxide and bilirubin production by tissue homogenates 315
Bis(2-chloroethyl)amine
Combined TLC–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 121
2,8-Bis(trifluoromethyl)-4-quinoline carboxylic acid
HPLC separation and extraction investigation for the simultaneous determination of mefloquine and its carboxylic acid metabolite 295
α-Bromoisovalerylurea
LC determination of diastereomeric glutathione conjugates and further derivatives of α-bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 67
Buflomedil
GC evaluation of buflomedil in biological samples employing the thermoionic specific detector 345
Caffeine
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277
Carbon monoxide
Correlation of carbon monoxide and bilirubin production by tissue honogenates 315
Carboxyphosphamide
Combined TLC–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 121
Catecholamines
Determination of catecholamines in body fluids 359
Catecholamines
Determination of catecholamines in body fluids 360
Cefixime
Application of HPLC of some antibiotics in clinical microbiology 257
Cephalexin
Analysis of cephalexin from canine skin biopsy by LC with UV–visible photodiode-array detection 103

Cephalexin
HPLC assay of cephalexin in serum and urine 162
Cephalosporin
Application of HPLC of some antibiotics in clinical microbiology 257
Ciprofloxacine
Application of HPLC of some antibiotics in clinical microbiology 257
Cyclodextrin glucosyltransferase
Biospecific sorption of cyclodextrin glucosyltransferase on physically modified starch 331
Cyclophosphamide
Combined TLC–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 121
Cyclosporin
Monitoring of cyclosporin in whole blood by reversed-phase LC on a butyl column 195
Dactinomycin
Reversed-phase HPLC of dactinomycin 166
3-Desacetylpancuronium
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 41
3-Desacetylpipecuronium
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 41
3-Desacetylvecuronium
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 41
Desipramine
Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative LC 287
Diazepam
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277
Dihydroxyphenylacetic acid
Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 9
D,L-*threo*-3,4-Dihydroxyphenylserine
Determination of D,L-*threo*-3,4-dihydroxyphenylserine and of the D- and L-enantiomers in human plasma and urine 219
Dopamine
Tritium isotope effect in reversed-phase HPLC analysis of dopamine and dihydroxyphenylacetic acid 9
Drugs
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277
Enzymes
Simple assay procedure for tyrosine hydroxylase activity by HPLC employing coulometric detection with minimal sample preparation 229
Enzymes
Biospecific sorption of cyclodextrin glucosyltransferase on physically modified starch 331
Eproxindine
Determination of eproxindine hydrochloride in human plasma by capillary GC 157
Erythromycin
HPLC of erythromycin propionyl ester and erythromycin base in biological fluids 93
Erythromycin propionyl ester
HPLC of erythromycin propionyl ester and erythromycin base in biological fluids 93
Ferritin receptor
Chromatographic detergent exchange in the preparation of membrane receptor protein 341
Flecainide
Rapid and inexpensive HPLC method for the quantification of flecainide in serum 351
Flunixin
GC analysis of flunixin in equine urine after extractive methylation 55
Flunoxaprofen
S-(+)-Flunoxaprofen chloride as chiral fluorescent reagent 131
Glafenic acid
Effective HPLC determination of glafenine in plasma: pharmacokinetic application 307
Glafenine
Effective HPLC determination of glafenine in plasma: pharmacokinetic application 307
Glutathione
LC determination of diastereomeric gluta-

thione conjugates and further derivatives of α-bromoisovalerylurea in rat bile and urine by electrochemically generated bromine 67
Haemoglobin
Separation of human haemoglobin alkylated at β93 cysteine from its native form by fast protein LC 335
2-Hydroxydesipramine
Isolation and identification of the glucuronide conjugate of 2-hydroxydesipramine by preparative LC 287
Hydroxy fatty acids
Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 1
Imipenem
Application of HPLC of some antibiotics in clinical microbiology 257
Insulin
Chromatographic analysis of the trinitrophenyl derivatives of insulin 239
Leukotrienes
Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 1
Leukotrienes
Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis – routine monitoring by HPLC 199
Lipoxins
Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 1
Mefloquine
HPLC separation and extraction investigation for the simultaneous determination of mefloquine and its carboxylic acid metabolite 295
Melperone
GC–MS study of urinary metabolism of melperone 144
Membrane proteins
Chromatographic detergent exchange in the preparation of membrane receptor protein 341
N-(2-Methyl-2-propyl)-3-oxo-4-aza-5α-androst-1-ene-17β-carboxamide
HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5α-androst-1-ene-17β-carboxamide, a 4-azasteroid, in human plasma from a phase I study 79
Mezlocillin
Application of HPLC of some antibiotics in clinical microbiology 257
Monohydroxyeicosatetraenoic acids
Reversed-phase ion-interaction chromatography of leukotrienes, lipoxins and related compounds 1
Monohydroxyeicosatetraenoic acids
Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severly burned patients and children with cystic fibrosis – routine monitoring by HPLC 199
Nalbuphine
Determination of nalbuphine by HPLC with electrochemical detection: application to clinical samples from post-operative patients 113
Nordiazepam
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277
Norfloxacine
Application of HPLC of some antibiotics in clinical microbiology 257
Octopamine
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 320
Ofloxacine
Application of HPLC of some antibiotics in clinical microbiology 257
Oxazepam
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277
4-Oxocyclophosphamide
Combined TLC–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 121
Pancuronium
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 41
Penems
Application of HPLC of some antibiotics in clinical microbiology 257
Penems

HPLC determination of the penem antibiotic (5*R*,6*S*)-2-aminomethyl-6-[(1*R*)-hydroxyethyl]-2-penem-3-carboxylic acid in human plasma and urine 269

Phencyclidine
Rapid confirmation of enzyme-multiplied immunoassay test (EMIT) for phencyclidine-positive urine samples with capillary GC–nitrogen–phosphorus detection 151

Phenethylamine
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 320

Phenobarbital
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277

Phenylethanolamine
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 320

Phosphoramide mustard
Combined TLC–photography–densitometry for the quantitation of cyclophosphamide and its four principal urinary metabolites 121

Pipecuronium
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 41

Piperacillin
Application of HPLC of some antibiotics in clinical microbiology 257

Prostaglandins
Arachidonic acid metabolites from polymorphonuclear leukocytes of healthy donors, severely burned patients and children with cystic fibrosis – routine monitoring by HPLC 199

Prostaglandins
Determination of prostaglandins and thromboxane in whole blood by HPLC with fluorimetric detection 209

Proteins
Chromatographic detergent exchange in the preparation of membrane receptor protein 341

Quinolones
Application of HPLC of some antibiotics in clinical microbiology 257

Serotonin
Automatic determination of serotonin in biological fluids by LC 326

Sotalol
Improved HPLC method for the analysis of serum sotalol 181

Steroids
HPLC determination of N-(2-methyl-2-propyl)-3-oxo-4-aza-5α-androst-1-ene-17β-carboxamide, a 4-azasteroid, in human plasma from a phase I study 79

Sulfamic acid diester
GC assay for the new antitumor agent sulfamic acid diester (NSC 329680) and its stability in buffer, blood and plasma 247

Temazepam
Use of micellar mobile phases and microbore column switching for the assay of drugs in physiological fluids 277

Thromboxane
Determination of prostaglandins and thromboxane in whole blood by HPLC with fluorimetric detection 209

Tocopherol
Two-dimensional HPLC system for the direct determination of tocopherol derivatives in plasma 29

Tyramine
Simultaneous determination of plasma phenethylamine, phenylethanolamine, tyramine and octopamine by HPLC using derivatization with fluorescamine 320

Tyrosine hydroxylase
Simple assay procedure for tyrosine hydroxylase activity by HPLC employing coulometric detection with minimal sample preparation 229

Vecuronium
Quantitation of pancuronium, 3-desacetylpancuronium, vecuronium, 3-desacetylvecuronium, pipecuronium and 3-desacetylpipecuronium in biological fluids by capillary GC using nitrogen-sensitive detection 41

Vitamin E
Two-dimensional HPLC system for the direct determination of tocopherol derivatives in plasma 29

Vitellogenin
Determination of vitellogenin in serum of rainbow trout (*Salmo gairdneri*) by high-performance gel permeation chromatography 138